

HyOctane

Forage Winter Triticale



HyOctane Forage Winter Triticale is a new awnletted variety that has been crossed between Wheat and Rye. This gives *HyOctane* the yield potential of rye grain and the feed quality of wheat. *HyOctane* can be used in many different applications such as silage, green chop, *hay, grazing or even cover cropping. *HyOctane* has an earlier heading date than most of the other Triticale's—making it a great choice for double cropping systems.



Seeding Rates:

Late Summer or Early Fall: 80-100 lbs/acre

Early to Mid Fall: 100-110 lbs/acre

Mid Fall: 110-120 lbs/acre

Planting Dates: August to November. *HyOctane* can also be late-spring planted for summer grazing, then cut the following year for forage after the plant has gone through the vernalization process.

Seed Yield: Although *HyOctane* is a Forage Winter Triticale, do not count it out for seed yield. *HyOctane* was a 130% of the check varieties in Minto and Manito-ba.

Harvesting for Feed: Best Feed Quality comes when *HyOctane* is in the late boot state. Protein can be found from 15-22%. At soft dough stage the Dry Matter Yield will typically double, but overall Feed Quality Protein will drop to 8-13%. The protein levels vary based on fertility and overall growing conditions.

Waste Management: *HyOctane* is a good choice for dairy waste management. It can consume as much as 250 units or more of nitrogen if the applications are applied uniformly. Always check nitrate levels before feeding.

Grazing Tips: Grazing fall triticale is one of the most overlooked forage producing attributes of spring planted winter triticale. If planted in the spring, *HyOctane* can realistically be grazed from spring through summer, and even into the fall. This does require proper management with fertility and some irrigation. If *HyOctane* does not go through a winter, it won't vernalize and go into the reproductive stage – leaving the plant vegetative until winter. This allows for better silage, green chop, and possibly even a May hay crop.

*For the hay producers: Please keep in mind that HyOctane is Awnletted not Awn less.

2 Year Trial Data Comparison

	Winter Survival Visual %	Dry Matter Yield (Kg/Ha)	Grain Yield (Kg/Ha)
HyOctane	97	14997.7	7316
Pika	98	13342.8	3117
Fridge	98	14785.0	5391
Bobcat	99	13443.0	4499
Check Mean	98	13856.9	4335.6
Sites*	3	2	3

*Taber, AB: 2007-08 Minto, MB: 2007-08 Taber, AB: 2006-07

Composition of Triticale Grain

Component	Percent of Dry Matter
Protein	19.71
Fiber	3.10
Fat	1.61
Calcium	.12
Phosphorus	.44
Total Sugars (as invert) Starch	5.74
	67.78

Source: Waibel et A., 1992, University of Minnesota

Bailey Seed & Grain, LLC

800-407-7713

www.baileyseed.com

4570 Ridge Dr. NE

Salem, OR 97301 tel:

503-362-9700 Fax:

503-362-1705